

1/13




Figure 1

2/13




Figure 2

3/13

a)

The viability of L1210 cells after exposure to converted material and free fatty acids

	Viability (%)
Medium control	99
Fatty acid/protein complexes:	
18:1:9c (HAMLET)	0
18:1:9tr	98
18:1:11c	1
18:1:6c	60
18:3:c	32
$\gamma$ 18:3:c	43
18:1:11tr	98
18:2:c	89
16:1:9c	73
16:1:9tr	98
20:1:11c	96
20:4:c	96
Fatty acids:	
18:1:9c	99
18:1:9tr	98
18:1:6c	97
18:1:11tr	96
18:2:c	98
18:3:c	99
$\gamma$ 18:3:c	99
16:1:9c	96
16:1:9tr	98
20:1:11c	99
20:4:c	98

a) Protein-lipid complexes




Figure 3

10/506903

4/13




Figure 4

5/13




Figure 5

6/13




Figure 6

10/506903

7/13

A)










Figure 7

10/506903

8/13



B)

Table I. The viability of L1210 cells after treatment with various forms of  $\alpha$ -lactalbumin.

	Cell viability (%)
Medium control	98
$\alpha$ -lactalbumin*:	
native	99
D87A	99
D87N	97
HAMLET†	4

\* the concentration was 1.0 mg/ml  
 † the concentration was 0.3 mg/ml


A)

Figure 8



9/13

A)



B)

Table II. Viability of L1210 cells after treatment with BAMLET and HAMLET.

Cell viability (%)	
Medium control	98
HAMLET	
0.2 mg/ml	67
0.3 mg/ml	9
BAMLET	
0.2 mg/ml	76
0.3 mg/ml	8

C)





Figure 9


10/506903

10/13

D)



E)



F)





Figure 9 con't



B)

Table III. The viability of L1210 cells after treatment with D87A- and D87N-BAMLET.

Cell viability (%)	
Medium control	98
$\alpha$ -lactalbumin*	
native	97
D87A-BAMLET	13
D87N-BAMLET	17
BAMLET†	0

\*the concentration was 0.5 mg/ml

†the concentration was 0.3 mg/ml

C)






Figure 10

12/13

D)



E)



F)




Figure 10 con't

10/506903

13/13




Figure 11